

---

# Loss of Reading and Central Vision Due to Macular Diseases - Therapeutic Management, Advances and Limitations

Gregg T. Kokame MD\*

Remarkable advances have recently developed in the management of macular diseases. There are now treatment options for diseases previously felt to be incurable, and for recovery of central vision for some of these patients. The macula is the central part of the retina (the nerve tissue lining the back of the eye). The macula provides for the fine and discriminating vision, needed for driving, reading and facial recognition. Thus, macular diseases result in loss of some of the most important visual tasks, ie. - the ability to read, the ability to drive, and the ability to recognize others. Vitreoretinal surgery is a rapidly changing subspecialty of ophthalmology, which has recently developed new approaches to macular diseases, such as macular holes, macular degeneration, and subretinal neovascularization. Research in Hawaii has contributed to these advances.

## Macular Holes

Age-related macular holes are holes in the central part of the retina, resulting in a missing spot in the center of vision and in distortion of vision (metamorphopsia) (Fig. 1). Macular holes are more common in females and increase in frequency with age, especially after age 55. Prevalence has been estimated to be 3.3 per 1000 in people over 55. This disease was previously felt to be untreatable until 1991, when the first pilot series of vitreous surgery showed closure of the macular hole, sealing of the retinal separation and improvement of central vision (Fig. 2).<sup>1</sup> Marked improvements in surgical techniques have resulted in a significant increase in success rate from 58% in the initial reported series to over 90% in recent series, including series reported from the Retina Center at Pali Momi. To take a disease previously untreatable six years ago to

a point of highly successful surgical management is truly a remarkable story in the field of vitreoretinal surgery.

The Retina Center at Pali Momi, (a division of Kapiolani Medical Center at Pali Momi), is one of 15 study centers around the nation participating in the randomized national multi-centered trials on macular hole surgery (The Vitrectomy for Prevention of Macular Hole Study<sup>2,3</sup> and The Vitrectomy for Macular Hole Study).<sup>4-6</sup> These studies have resulted in significant contributions to the literature and represent the only well-controlled randomized studies on macular hole surgery. The Retina Center at Pali Momi is studying the use of serum growth factors to stimulate higher macular hole closure rate and vitrectomy with an intraocular gas bubble to prevent macular hole development in early stages of this disease.<sup>3</sup> In previously published studies ultrasonographic techniques were utilized to demonstrate tangential traction within the vitreous gel causing macular holes<sup>7</sup>, and macular hole surgery was shown to allow patients even with chronic macular holes of many years in duration to recover useful central vision following surgery.<sup>8</sup>

## Age-Related Macular Degeneration

Age-related macular degeneration (AMD) is a degenerative disease of the central retina, which results in a gradual scarring of support tissues. AMD is the most common causes of reportable blindness in the United States, as well as here in Hawaii. It is most common in Caucasians, especially fair and blue-eyed individuals, and less common, although not infrequently seen in Asians. Presently, preventative treatment is not proven, but studies on mineral, antioxidant, and vitamins supplements are ongoing. Laser treatment in the early stages of the disease is being studied in randomized trials.

AMD has two categories - the dry form and the wet form. The mainstay of treatment is laser surgery for the wet form which is the most common cause of severe vision loss. In the dry form there is gradual scarring with slow and gradual vision loss. In the wet form, choroidal neovascular membranes (CNVM) rapidly proliferate causing bleeding, scarring and leaking and rapid vision loss (Fig. 3). Laser treatment to obliterate the CNVM has been shown in large national multi-centered trials to decrease the risk of further vision loss.<sup>10-11</sup> A fluorescein angiogram (office procedure) is used to visualize the extent of CNVM (Fig. 3). Fluorescein dye is injected into an arm vein and an ophthalmic fundus camera with specific filters for the fluorescent light visualizes the dye in the retinal vessels.

\*Clinical Professor of Surgery  
Division of Ophthalmology  
Department of Surgery  
University of Hawaii School of Medicine  
Medical Director, The Retina Center at  
Pali Momi, Kapiolani Medical Center  
at Pali Momi, Aiea, Hawaii.

Reprint requests to:  
Gregg T. Kokame, MD  
321 N. Kuakini Street  
Suite 307  
Honolulu, Hawaii 96817

Fig 1.—Photograph of macular hole in left eye. Note visible hole approximately 800  $\mu$ m in size in center of macula. Visual acuity is 20/200.



Fig. 2.—Postoperative photograph following macular hole surgery. Note that macular hole has sealed. Visual acuity is 20/40.




Fig. 3.—Fluorescein angiogram of retinal pigment epithelial detachment (RPED) or blister-like formation in pigment epithelium due to subretinal neovascularization and AMD. Note bright spot at superonasal edge of blister consistent with SRNVM, but partially obscured by fluorescence within the RPED.




Fig. 4.—ICG angiogram demonstrating bright, focal area of hyperfluorescence consistent with SRNVM in same eye as in Figure 3. Note absence of fluorescence obscuring SRNVM within the RPED.



Fig. 5.—Photograph of SRNVM in patient with idiopathic inflammatory disease of retina, called punctate inner choroidopathy. Note the grey subretinal lesion extending into central fovea, as well as serous retinal detachment. Visual acuity is 20/200.




Fig. 6.—Postoperative photograph following subretinal surgery to remove SRNVM showing focal hypopigmented scar in temporal macula. Note absence of grey membrane and serous retinal detachment. Visual acuity has improved to 20/30.



The images are captured on film or on digital camera with computer interface, and laser treatment can be guided to the area of CNVM. Unfortunately, the percentage of patients with the wet form treatable with fluorescein angiographic criteria is less than 15%. A new imaging technology utilizing indocyanine green (ICG) may allow more precise localization of CNVM in select cases (Fig. 4). This imaging technology uses digital imaging, because of the higher sensitivity of digital cameras or video cameras to the fluorescence of the ICG dye in the infrared range. This imaging technique has become available in Hawaii for just over a year. With further experience and research with this modality, more patients may become eligible for laser surgery, although we have often found the two imaging modalities are complimentary. For patients not treatable with laser surgery, the prognosis for central vision is poor, although the majority of patients will have normal peripheral vision. Ongoing studies are evaluating medical treatments of CNVM by anti-angiogenic modalities, such as oral thalidomide, vascular endothelial growth factor antibodies, and radiation therapy.

Choroidal neovascular membranes (CNVM) are abnormal blood vessels growing into the subretinal space through retinal pigment epithelium. The most common disease is AMD, but there are other entities, which cause retinal scars and CNVM ingrowth. These include ocular histoplasmosis syndrome (a scarring disease associated with the fungus, *Histoplasma capsulatum*, common in the midwest), high myopia, angioid streaks (associated most commonly with pseudoxanthoma elasticum), trauma with choroidal ruptures, and multiple viral or idiopathic inflammatory diseases of the retina. Because CNVM is subretinal under the nerve tissue thus limiting surgical access, surgical removal was not considered until the first successful surgical series of CNVM removal through a small retinotomy in 1991.<sup>12</sup> Subretinal surgery has since developed remarkably with improved techniques and instrumentation. Most promising visual results have been in patients with ocular histoplasmosis or idiopathic causes (Figs 5-6).<sup>13</sup> The initial results of surgical removal of CNVM in AMD were disappointing, but research is ongoing, including exciting possible new treatment avenues of neuroretinal and retinal pigment epithelial cell transplantation. Subretinal surgery to remove hemorrhage using tissue plasminogen activator has successfully restored vision in subretinal hemorrhage due to retinal macroaneurysms and AMD.

Macular diseases cause loss of some of the most important visual abilities. Recent diagnostic, therapeutic and surgical advances have resulted in recovery of central vision for many patients with these diseases. Ongoing research has led to advances in this rapidly changing field, much occurring within this decade.

## References

1. Kelly NE, Wendel RT. Vitreous surgery for idiopathic macular holes. Results of a pilot study. *Arch Ophthalmol* 1991; 109:654 - 659.
2. de Bustros S, The Vitrectomy for Prevention of Macular Hole Study Group. Vitrectomy for prevention of macular holes. Results of a randomized multicenter clinical trial. *Ophthalmology* 1994; 101: 1055 - 1060. (Kokame, GT - Principal Investigator, Honolulu Center).
3. Kokame GT, de Bustros S, The Vitrectomy for Prevention of Macular Hole Study Group. Visual acuity as a prognostic indicator in stage I macular holes. *Am J Ophthalmol* 1995; 120: 112 - 114.
4. Kim JW, Freeman WR, El-Haig W, Maguire AM, Arevalo JF, Azen SP, The Vitrectomy for Macular Hole Study Group. Baseline characteristics, natural history and risk factors to progression in eyes with stage II macular hole. Results from a randomized prospective clinical trial. *Ophthalmology* 1995; 102: 1818 - 1827. (Kokame GT - Principal Investigator, Honolulu Center).
5. Kim JW, Freeman WR, Azen AP, El-Haig W, Klein DJ, Bailey IL, The Vitrectomy for Macular Hole Study Group. Prospective randomized trial of vitrectomy or observation for stage II macular holes. *Am J Ophthalmol* 1996; 121: 605 - 614. (Kokame GT - Principal Investigator, Honolulu Center).
6. Freeman WR, Jung KW, Azen SP, El-Haig W, Mishell D, Bailey IL, The Vitrectomy for Macular Hole Study Group. Vitrectomy for the treatment of full-thickness stage III or IV macular holes: results of a randomized clinical trial. *Arch Ophthalmol* 1997; 115: 11-21. (Kokame GT - Principal Investigator, Honolulu Center).
7. Kokame GT. Clinical correlation of ultrasonographic findings in macular holes. *Am J Ophthalmol* 1995; 119: 441 - 451.
8. Kokame GT. Macular hole surgery for chronic macular holes. *Retina* 1996; 16: 75 - 78.
9. Kokame GT. Early stage of macular hole in a severely myopic eye. *Am J Ophthalmol* 1995; 119: 240 - 242.
10. Macular Photocoagulation Study Group: Argon laser photocoagulation for neovascular maculopathy after five years: results from randomized trials. *Arch Ophthalmol* 1991; 109: 1109 - 1114.
11. Macular Photocoagulation Study Group: Krypton laser photocoagulation for neovascular lesions of age-related macular degeneration: results of a randomized clinical trial. *Arch Ophthalmol* 1990; 108: 816 - 824.
12. Thomas MA, Kaplan HJ. Surgical removal of subfoveal neovascularization in the presumed ocular histoplasmosis syndrome. *Am J Ophthalmol* 1991; 111: 1-7.
13. Gass JDM. Biomicroscopic and histological considerations regarding the feasibility of surgical excision of subfoveal neovascular membranes. *Am J Ophthalmol* 1994; 118: 285 - 298.

## The Mama Lau Cancer Care Center

(opening in October 1997)

Providing leadership in cancer care services  
and technologies for the people of Hawaii.

Coming in early 1998...  
Gamma Knife technology...




photo of Gamma Knife equipment

Gamma Knife is a tremendous alternative to brain surgery. The first in Hawaii, this service will provide non-invasive, relatively painless, radiosurgical treatment of brain tumors and lesions.



**St. Francis Medical Center**

*A subsidiary of St. Francis Healthcare System of Hawaii*

2230 Liliha Street  
Honolulu • Hawaii  
(808) 547-6011

### Gamma Knife Benefits

- Superior outcomes
- Reduced health risks
- No general anesthesia
- No incision
- Overnight hospitalization
- Significantly reduced costs